

54% OF NEW HIV DIAGNOSES



35

Average age

are made in men who have sex with men (MSM)

38,4%

of MSM showed a late diagnosis (<350 CD4**)

1

in 3

new diagnoses are made in individuals < 30

HIV

In 2015, 1,837 new cases of HIV in MSM were diagnosed in Spain

1

in 10

MSM who show risk behaviours are infected with HIV

1 in 3

cases of syphilis and gonorrhoea in MSM were transmitted through oral sex

1 in 3

cases of syphilis


in MSM were also diagnosed with HIV

1 in 5

cases of gonorrhoea

In recent years, there has been a significant increase in sexually transmitted infections (STI)

>50% of STIs are diagnosed in MSM



● Gonorrhoea ● Syphilis

SOURCE: NATIONAL EPIDEMIOLOGICAL MONITORING NETWORK

NO CONDOM
no turning back.



Source: Spanish National AIDS Plan
Further information:
<http://www.msc.es/ciudadanos/enf/infecciones/enf/transmisibles/sida/home.htm>
<http://www.msc.es/ciudadanos/enf/infecciones/enf/transmisibles/sida/homelT5.htm>

*Trends (2009-2015): Stable
• The rate of new diagnoses of HIV in MSM has not gone down in recent years.
• The percentage of cases with late diagnoses has not gone down in recent years.

**A type of cells that are an essential part of the immune system. Their main function is to activate the system by warning it of the presence of pathogens or of an erroneous replication of human cells, so it can deal with them and correct the situation.